arXiv:1212.5883v4 [math.NT] 22 Jan 2013

ON A GENERALIZATION OF DEURING'S RESULTS

KEN-ICHI SUGIYAMA

Department of Mathematics and Informatics, Faculty of Science,
Chiba University, 1-33 Yayoi-cho Inage-ku,
Chiba 263-8522, Japan
e-mail address : sugiyama@math.s.chiba-u.ac.jp

Abstract. Using the Dieudonné theory we will study a reduction of an abelian variety with complex multiplication. Our results may be regarded as a generalization of a classical theorem due to Deuring for a CM-elliptic curve. We will also discuss a sufficient condition for a prime at which the reduction of a CM will be maximal.

Key words: complex multiplication, a maximal curve over a finite field.
AMS classification 2010: 11G10, 11G15, 11G20, 11G30, 14F30, 14F40, 14G50, 14H40.

1. Introduction

Let E be an elliptic curve over the field of rational numbers $\mathbb{Q}$ with complex multiplication (which will be abbreviated by *CM*) by the integer ring of an imaginary quadratic field K. Suppose that it has a non-singular reduction E_p at a prime p. The classical Deuring's theorem says that it is ordinary or supersingular if p completely splits or remains prime in K, respectively. Here we note that E_p is ordinary (resp. supersingular) if and only if the p-divisible group $E_p[p^\infty]$ is $\mathbb{L}$ (resp. $G_{1,1}$) (see \$2 for details). Moreover if E_p is super-singular and if $p \geq 5$ the characteristic polynomial of p-power Frobenius on an l-adic Tate module ($l \neq p$) is $t^2 + p$ and the number of $\mathbb{F}_{p^2}$-rational points attains the Hasse-Weil upper bound:

$$|E_p(\mathbb{F}_{p^2})| = 1 + p^2 + 2p.$$

In this report we will generalize these results to a proper smooth curve over $\mathbb{Q}$ of higher genus with CM.

We first explain notation which will be used. An algebraic closure of a field F is written by $\overline{F}$. For a scheme X over F its base change to an field extension F' of F is denoted by $X \otimes_F F'$. For simplicity we sometimes write $\overline{X} := X \otimes_F \overline{F}$. Suppose that F is a finite field of q-elements and let V be a proper smooth variety. Then the characteristic polynomial of the q-power Frobenius on the first l-adic étale cohomology group $H^1_{et}(\overline{V}, \mathbb{Q}_l)$ ($l \neq \mathrm{char}(F)$) is described by $\Phi_{V,q}(t)$.

Let k be a finite field of characteristic $p \geq 5$. An abelian variety J of dimension g over k will be called *supersingular* (resp. *superspecial*) if $\overline{J}$ is isogeneous (resp. isomorphic) to a product of supersingular elliptic curves. Moreover it is defined to be *ordinary* if the group of p-torsion points $\overline{J}[p]$ is isomorphic to $(\mathbb{Z}/p\mathbb{Z})^g$. Let $W(k)$ be the ring of Witt vectors whose coefficients in k. An abelian scheme $\mathcal{A}$ over $W(k)$ of relative dimension g will be called *with CM* (or sometimes *with CM by $\mathcal{O}_K$*) if the endomorphism ring over $W(\overline{k})$ contains the integer ring $\mathcal{O}_K$ of a CM-field K satisfying $[K:\mathbb{Q}] = 2g$. Let K_0 be the totally real subfield with $[K_0:\mathbb{Q}] = g$. We assume that p is unramified in K and let

$$p = \mathfrak{P}_1 \cdots \mathfrak{P}_t \tag{1}$$

be the prime factorization in K_0.

Theorem 1.1. *Suppose that every $\mathfrak{B}_i$ remains prime in K. Then the special fiber A is supersingular. Moreover if $t = g$ (i.e. p completely splits in K_0) it is superspecial.*

This result gives us an information of the characteristic polynomial of a Frobenius. In fact suppose that the assumption is satisfied. Then **Theorem 1.1** says that there is a positive integer m so that $A \otimes_k \mathbb{F}_{p^m}$ is isogeneous to a product of elliptic curves $\{E_i\}_{1\leq i\leq g}$ defined over $\mathbb{F}_{p^m}$ and $\Phi_{A,p^m}(t) = \prod_{i=1}^{g} \Phi_{E_i,p^m}(t)$. Since $p \geq 5$, $\Phi_{E_i,p^m}(t)$ is a one of the following([12] **Theorem 4.1**):

$$t^2 + p^m, \quad t^2 \pm p^{\frac{m}{2}} t + p^m, \quad t^2 \pm 2p^{\frac{m}{2}} t + p^m,$$

where the last two occur only when m is even.

Theorem 1.2. *Suppose that p completely splits in K. Then A is ordinary.*

In **Theorem 1.1** if $t = g$ and $k = \mathbb{F}_p$, we will obtain a stronger result.

Theorem 1.3. *Let $\mathcal{A}$ be an abelian scheme over $\mathbb{Z}_p$ with CM. Suppose that p completely splits in K_0:*

$$p = \mathfrak{P}_1 \cdots \mathfrak{P}_g,$$

and that every $\mathfrak{P}_i$ remains prime in K. Then A is a product of supersingular elliptic curve $\{E_i\}_{1\leq i\leq g}$,

$$A = E_1 \times \cdots \times E_g,$$

over $\mathbb{F}_p$. In particular $\Phi_{A,p}(t) = (t^2 + p)^g$.

A proper smooth curve C defined over a number field F of genus g is called *a CM-curve* if the endomorphism ring of the Jacobian variety $\mathrm{Jac}(\overline{C})$ contains $\mathcal{O}_K$, where K is a CM-field satisfying $[K:\mathbb{Q}] = 2g$. We say a finite prime v of F is *good* if the reduction C_v is nonsingular and moreover mention it *ordinary* (resp. *supersingular*, *superspecial*) if so is $\mathrm{Jac}(C_v)$. **Theorem 1.1**, **Theorem 1.2**, **Theorem 1.3** and the Hasse-Weil formula yield the following consequence, which generalizes the Deuring's result which has been explained before.

Theorem 1.4. *Let C be a proper smooth curve of genus g over $\mathbb{Q}$ with CM and $p(\geq 5)$ a good prime. Then we have the followings:*

(1) *If p completely splits in K, C_p is ordinary.*

(2) *Let*

$$p = \mathfrak{P}_1 \cdots \mathfrak{P}_t,$$

be the prime factorization in K_0. *If every* $\mathfrak{P}_i$ *remains prime in* K, C_p *is supersingular.*

(3) *Suppose that* $t = g$ *in the above (2) (i.e. p completely splits in* K_0*). Then* $\mathrm{Jac}(C_p)$ *is a product of supersingular elliptic curves over* $\mathbb{F}_p$ *and* $|C(\mathbb{F}_{p^2})|$ *attains the Hasse-Weil upper bound:*

$$|C(\mathbb{F}_{p^2})| = 1 + p^2 + 2gp.$$

Let C and p be as in **Theorem 1.4**.

Corollary 1.1. *Suppose that* K *is a cyclotomic field* $\mathbb{Q}(\zeta_N)$ *where* ζ_N *is a primitive* N*-th root of unity that satisfies* $\phi(N) = 2g$, *where* ϕ *is the Euler function.*

(1) *Suppose that* $p \equiv 1(N)$. *Then* C_p *is ordinary.*
(2) *Suppose there is a positive integer* h *so that* $p^h \equiv -1(N)$. *Then* C_p *is supersingular.*
(3) *Suppose that* $p \equiv -1(N)$. *Then* $\mathrm{Jac}(C_p)$ *is a product of supersingular elliptic curves over* $\mathbb{F}_p$ *and*

$$|C(\mathbb{F}_{p^2})| = 1 + p^2 + 2gp.$$

Corollary 1.2. *Suppose that* $K = \mathbb{Q}(\zeta_M + \zeta_M^{-1}, \zeta_d)$, *where* M *is a positive integer satisfying* $\phi(M) = 2g$ *and* $d = 3$, *or* 4.

(1) *Suppose that* $p \equiv 1(d)$ *and* $p \equiv \pm 1(M)$. *Then* C_p *is ordinary.*
(2) *Suppose that* $p \equiv -1(d)$. *Then* C_p *is supersingular.*
(3) *Suppose that* $p \equiv -1(d)$ *and that* $p \equiv \pm 1(M)$. *Then* $\mathrm{Jac}(C_p)$ *is a product of supersingular elliptic curves over* $\mathbb{F}_p$ *and*

$$|C(\mathbb{F}_{p^2})| = 1 + p^2 + 2gp.$$

In the final section we will show some examples of curves over $\mathbb{F}_p$ that attain the Hasse-Weil upper bound. We hope that our theorems may offer a new construction of such a curve.

Let us briefly explain how the theorems will be proved. For simplicity we assume that p completely splits in K_0: $p = \mathfrak{P}_1 \cdots \mathfrak{P}_g$. Using Dieudonné-Manin's theory and Grothendieck's theorem we will show that p-divisible group $A[p^\infty]$ is a product

$$A[p^\infty] = G_1 \times \cdots \times G_g, \tag{2}$$

whose every component has height 2. Moreover if all $\mathfrak{P}_i$'s completely split in K we will prove that $G_i \simeq \mathbb{L}$ $(\forall i)$ and A is ordinary. On the other hand if all of them remain prime in K it will be shown that $G_i \simeq G_{1,1}$ $(\forall i)$ and the a-number of A is g. Then using a theorem due to Oort we will conclude that A is a product of supersingular elliptic curves over $\overline{\mathbb{F}}_p$:

$$A = E_1 \times \cdots \times E_g. \tag{3}$$

Thus we obtain **Theorem 1.2** and the last half of **Theorem 1.1**. In order to prove **Theorem 1.3** we have to show that (3) is defined over the prime field. We will achieve this by establishing that (2) is defined over $\mathbb{F}_p$ with a help of Tate's theorem.

Let us mention precedent results of a generalization of Deuring's results. Using the Kraft's diagrams instead of the Dieudonné module, Zaytsev has completely determined p-torsion group of an abelian variety with dimension less than 4 [13]. Our **Theorem 1.1** and **Theorem 1.2** are contained in his results if $\dim A \leq 3$.

Acknowledgement. The author appreciates Professor A. Zaytsev for his interest and useful remarks. This research is partially supported by JSPS grants Kiban(C)22540068.

2. A REVIEW OF p-DIVISIBLE GROUPS.

In the present section we will summarize facts of p-divisible groups and the Dieudonné functor which will be used later. The references are [2], [4], [6], [7] and [8]. Throughout the section k will be a field of characteristic p.

Definition 2.1. *Let h be a nonnegative integer.* A p-divisible group G of height $h(G) = h$ *over k is an inductive system of finite group schemes $G_i \to \operatorname{Spec} k$, $(i \geq 1)$, satisfying*

1. *the dimension of the coordinate ring $k[G_i]$ over k equals to $p^{h \cdot i}$,*
2. *p^i annihilates G_i,*
3. *there are inclusions $G_i \hookrightarrow G_{i+1}$ such that*
$$G_{i+1}[p^i] = G_i.$$

Here we have used the standard notation
$$X[f] = \operatorname{Ker}[X \xrightarrow{f} X],$$
for an endomorphism f of a group scheme X. Note that by definition $G_{i+j}/G_i = G_j$. We will write $G = \lim_{\to} G_i$. The basic examples are $\mathbb{Q}_p/\mathbb{Z}_p := \lim_{\to} \mathbb{Z}/p^n\mathbb{Z}$ and $\mathbb{G}_m[p^\infty] := \lim_{\to} \mathbb{G}_m[p^n]$, which have height 1. Let A be an abelian variety over k of dimension g. Then $A[p^\infty] := \lim_{\to} A[p^n]$ is a p-divisible group of height $2g$. The collection of Cartier dual $\{G_i^\vee\}_i$ of a p-divisible group $G = \lim_{\to} G_i$ is also a p-divisible group. We denote it by $G^\vee$ and call it *the Serre dual*. Taking the Serre dual induces an involution of the category of p-divisible groups. Note that $\mathbb{Q}_p/\mathbb{Z}_p$ and $\mathbb{G}_m[p^\infty]$ are Serre dual to each other.

Let $W(k)$ be the ring of Witt vectors whose coefficients are in k and consider a non-commutative ring $\mathcal{D}(k) = W(k)[F, V]$ generated by semi-linear operators F and V with relations
$$FV = VF = p, \quad F\lambda = \lambda^\sigma F, \quad \lambda V = V\lambda^\sigma, \quad \lambda \in W(k).$$
Here σ is the lift of p-power Frobenius automorphism of k. There is a contra-variant functor $\mathbb{M}$ called *the Dieudonné functor* from an abelian category of finite group schemes over k of p-power order to one of left $\mathcal{D}(k)$-modules of finite length. (Here *the order* $|\Gamma|$ of a finite group scheme Γ is defined to be the dimension of its coordinate ring over k). It is known that the length of $\mathbb{M}(\Gamma)$ is equal to $\log_p |\Gamma|$. Due to Dieudonné, Cartier, Barsotti and Oda,

it gives an anti-equivalence of these categories ([7] **Theorem 4**). Let $G = \lim_{\rightarrow} G_i$ be a p-divisible group. We define *the Dieudonné module* as $\mathbb{M}(G) := \lim_{\leftarrow} \mathbb{M}(G_i)$ which is a $\mathcal{D}(k)$-module by definition. It is a free $W(k)$-module whose rank is $h(G)$.

A homomorphism of p-divisible groups whose kernel is finite will be called *an isogeny.* Let X and Y are p-divisible group. If there is an isogeny between them we say X and Y are *isogeneous* and write $X \sim Y$. This notion defines an equivalence relation on the set of p-divisible groups.

Fact 2.1. [6] *Suppose that k is algebraically closed and let G be a p-divisible group over k. Then there is an isogeny:*

$$G \sim \prod_i G_{d_i,c_i}.$$

Temporary we assume that k is an algebraic closed field. A component $G_{d,c}$ is parametrized by two non-negative coprime integers d and c. We call d *the dimension* of G and c is defined by $c = h - d$. By definition $G_{c,d} = (G_{d,c})^\vee$ and it is known that $G_{d,c}$ is characterized by the height h and *the slope* $d/h = d/(c+d)$. A p-divisible group over k will be mentioned *simple* if it is isomorphic to $G_{d,c}$ for a certain coprime pair (c, d). A simple p-divisible group G is isomorphic to $G_{d,c}$ if and only if there is a positive integer (m, n) so that

$$G[F^m] = G[V^n], \quad \frac{n}{m+n} = \frac{d}{c+d}, \tag{4}$$

which follows from ([2] Chapter IV):

$$\mathbb{M}(G_{d,c}) = \mathcal{D}(k)/(F^c - V^d).$$

In fact for a simple p-divisible group G the above equation shows that $G = G_{d,c}$ if and only if

$$\mathbb{M}(G)/V^n\mathbb{M}(G) = \mathbb{M}(G)/F^m\mathbb{M}(G),$$

for a pair of positive integers (m, n) satisfying $d/(c + d) = n/(m + n)$, which is equivalent to (4). The fundamental examples are $G_{1.0} = \mathbb{G}_m[p^\infty]$, $G_{0.1} = \mathbb{Q}_p/\mathbb{Z}_p$ and $G_{1.1} \simeq E[p^\infty]$ where E is an supersingular elliptic curve. Here is a classification of p-divisible groups whose height is less than 3 ([2], p.93):

1. $h(G) = 0$ iff $G = 0$.
2. If $h(G) = 1$, then $G = G_{1.0}$, or $G_{0.1}$.
3. If $h(G) = 2$, G is the one of the followings,

$$G_{1.0}^2, \quad G_{1.0} \times G_{0.1}, \quad G_{0.1}^2, \quad G_{1.1}.$$

We will write $\mathbb{L} = G_{1.0} \times G_{0.1}$.

Let A be an abelian variety over k of dimension g. We define *p-rank* $f(A) = f$ as an integer such that $A[p](\overline{k}) \simeq (\mathbb{Z}/p\mathbb{Z})^f$, which equals to $\dim_{\mathbb{F}_p} \mathrm{Hom}(\mathbb{G}_m[p], \overline{A}[p])$. Let α_p be a finite group scheme defined by $\alpha_p := \mathrm{Spec}\overline{k}[x]/(x^p)$. Then we set $a(A) = \dim_{\overline{k}} \mathrm{Hom}(\alpha_p, \overline{A}[p])$

and call it the *a-number*. It is known that $0 \leq f(A) \leq g$, and $0 \leq a(A) + f(A) \leq g$. We say A is *ordinary* if $f(A) = g$.

Fact 2.2. [1] *Let A be an abelian variety over k of dimension g. Then the following are equivalent:*

1. *A is ordinary,*
2. *$\overline{A}[p^\infty] \sim \mathbb{L}^g$.*

This is well-known if $g = 1$. Let E be an elliptic curve defined over an algebraic closed field of characteristic p. $E[p^\infty]$ has height two and is isomorphic to $\mathbb{L}$ or $G_{1.1}$ according that it is ordinary or supersingular, respectively. Thus E is supersingular if and only if $E[p^\infty] = G_{1.1}$. Since there is an exact sequence

$$0 \to \alpha_p \to G_{1.1}[p] \to \alpha_p \to 0, \tag{5}$$

this is also equivalent to $a(E) = 1$. These are generalized to a higher dimensional abelian varieties. The following fact is due to Deligne, Oort, Shioda and Tate.

Fact 2.3. *Let A an abelian variety over an algebraic closed field k with dimension g. Suppose that*

$$A[p^\infty] \sim G_{1.1}^g.$$

Then A is super-singular.

Fact 2.4. *(*[9] **Theorem 2***) Let A an abelian variety over an algebraic closed field k of with dimension g. Suppose that*

$$\alpha(A) = g.$$

Then A is superspecial.

For an abelian variety A, following Milne and Waterhaus[7], we set $T_p(A) := \mathbb{M}(A[p^\infty])$.

Fact 2.5. [7] **Theorem 6**. *Let A and B are abelian varieties over a finite field k. Then*

$$\mathrm{Hom}_k(A, B) \otimes \mathbb{Z}_p \simeq \mathrm{Hom}_{\mathcal{D}(k)}(T_p(B), T_p(A)).$$

Fact 2.6. [4] *Let A be the special fiber of an abelian scheme $\mathcal{A}$ over $W(k)$. Then $T_p(A)$ is isomorphic to the first de Rham cohomology group $H^1_{\mathrm{DR}}(\mathcal{A}/W(k))$ as $\mathcal{D}(k)$-modules.*

3. A REDUCTION OF AN ABELIAN VARIETY WITH CM

Let $\mathcal{A}$ be an abelian scheme over $W(\overline{\mathbb{F}}_p)$ with CM by $\mathcal{O}_K$. Then $H^1_{DR}(\mathcal{A}/W(\overline{\mathbb{F}}_p))$ is isomorphic to $\mathcal{O}_K \otimes_{\mathbb{Z}} W(\overline{\mathbb{F}}_p)$. We assume that p is unramified in K and let

$$p = \mathfrak{P}_1 \cdots \mathfrak{P}_t,$$

be a prime factorization in K_0 so that $\mathfrak{P}_i$ remains prime (resp. completely splits) in K for $1 \leq i \leq s$ (resp. $s+1 \leq i \leq t$). Thus we have a prime factorization in K:

$$p = \tilde{\mathfrak{P}}_1 \cdots \tilde{\mathfrak{P}}_s \tilde{\mathfrak{P}}_{s+1} \tilde{\mathfrak{P}}'_{s+1} \cdots \tilde{\mathfrak{P}}_t \tilde{\mathfrak{P}}'_t \tag{6}$$

Here "′" means the complex conjugation. We set $\mathcal{P}_{inert} := \{\mathfrak{P}_1, \cdots, \mathfrak{P}_s\}$ and $\mathcal{P}^{f=1}_{inert} := \{\mathfrak{P} \in \mathcal{P}_{inert} \,|\, f(\mathfrak{P}/p) = 1\}$, where $f(\mathfrak{P}/p)$ is the inertia degree. Similarly $\mathcal{P}_{split} :=$

$\{\mathfrak{P}_{s+1},\cdots,\mathfrak{P}_t\}$ and $\mathcal{P}_{split}^{f=1} := \{\mathfrak{P}\in\mathcal{P}_{split}\,|\,f(\mathfrak{P}/p)=1\}$. Note that $\mathcal{O}_K\otimes_{\mathbb{Z}}\mathbb{Z}_p$ acts on the p-divisible group $A[p^\infty]$ of the special fiber A. According to (6) we have a decomposition

$$\mathcal{O}_K\otimes_{\mathbb{Z}}\mathbb{Z}_p \simeq W(\mathbb{F}_{\tilde{\mathfrak{P}}_1})\times\cdots\times W(\mathbb{F}_{\tilde{\mathfrak{P}}_s})\times\{W(\mathbb{F}_{\tilde{\mathfrak{P}}_{s+1}})\times W(\mathbb{F}_{\tilde{\mathfrak{P}}'_{s+1}})\}\times\cdots\times\{W(\mathbb{F}_{\tilde{\mathfrak{P}}_t})\times W(\mathbb{F}_{\tilde{\mathfrak{P}}'_t})\},$$

where $\mathbb{F}_{\tilde{\mathfrak{P}}}$ is the residue field. Using this we define e_i to be an element of $\mathcal{O}_K\otimes_{\mathbb{Z}}\mathbb{Z}_p$ corresponding to $(0,\cdots,0,1,0,\cdots,0)$ where i places in the i-th position from the left. Set $G_i := e_iA[p^\infty]$ and $\Gamma_j := G_{s+(2j-1)}\times G_{s+2j}$ $(1\le j\le t-s)$. Then,

$$A[p^\infty] = G_1\times\cdots\times G_s\times\Gamma_1\times\cdots\times\Gamma_{t-s}. \tag{7}$$

and we obtain,

$$\mathbb{M}(G_i) = e_iT_p(A) = e_iH^1_{DR}(\mathcal{A}/W(\overline{\mathbb{F}}_p)) = e_i(\mathcal{O}_K\otimes_{\mathbb{Z}}W(\overline{\mathbb{F}}_p)),$$

Here the second identity is due to **Fact 2.6**. For a prime factor $\tilde{\mathfrak{P}}$ of p in K we denote the corresponding factor of $A[p^\infty]$ by $G_{\tilde{\mathfrak{P}}}$. Let $\mathfrak{P}$ be a prime of K_0 dividing p. Then we define p-divisible subgroup $G_{\mathfrak{P}}$ of $A[p^\infty]$ as follows. If $\mathfrak{P}$ is contained in $\mathcal{P}_{inert}$ define $G_{\mathfrak{P}} := G_{\tilde{\mathfrak{P}}}$ where $\tilde{\mathfrak{P}}$ is the unique prime of K over $\mathfrak{P}$. Since $\mathbb{F}_{\tilde{\mathfrak{P}}}\simeq\mathbb{F}_{p^{2f(\mathfrak{P}/p)}}$,

$$h(G_{\mathfrak{P}}) = \mathrm{rank}_{W(\overline{\mathbb{F}}_p)}\mathbb{M}(G_{\tilde{\mathfrak{P}}}) = 2f(\mathfrak{P}/p). \tag{8}$$

On the other hand if it splits: $\mathfrak{P} = \tilde{\mathfrak{P}}\times\tilde{\mathfrak{P}}'$, we define $G_{\mathfrak{P}} := G_{\tilde{\mathfrak{P}}}\times G_{\tilde{\mathfrak{P}}'}$. Since $\mathbb{F}_{\tilde{\mathfrak{P}}}\simeq\mathbb{F}_{\tilde{\mathfrak{P}}'}\simeq\mathbb{F}_{p^{f(\mathfrak{P}/p)}}$, a similar observation shows

$$h(G_{\tilde{\mathfrak{P}}}) = h(G_{\tilde{\mathfrak{P}}'}) = f(\mathfrak{P}/p),\quad h(G_{\mathfrak{P}}) = h(G_{\tilde{\mathfrak{P}}}) + h(G_{\tilde{\mathfrak{P}}'}) = 2f(\mathfrak{P}/p). \tag{9}$$

Lemma 3.1. *For $\mathfrak{P}\in\mathcal{P}_{split}^{f=1}$, $G_{\mathfrak{P}}$ is isomorphic to $\mathbb{L}$.*

Proof. By (9) $h(G_{\tilde{\mathfrak{P}}}) = h(G_{\tilde{\mathfrak{P}}'}) = 1$ and the classification of p-divisible groups implies that $G_{\tilde{\mathfrak{P}}}$ or $G_{\tilde{\mathfrak{P}}'}$ is one of $\{G_{1.0}, G_{0.1}\}$. Notice that $G_{1.0}$ (resp. $G_{0.1}$) is characterized by the fact V (resp. F) is an isomorphism on $\mathbb{M}(G_{1.0})$ (resp. $\mathbb{M}(G_{0.1})$). Take a positive integer r so that $\pi = F^r$ is contained in $\mathcal{O}_K$ and let

$$(\pi) = \tilde{\mathfrak{P}}^a(\tilde{\mathfrak{P}}')^{a'}\delta,\quad (\delta,\tilde{\mathfrak{P}}) = (\delta,\tilde{\mathfrak{P}}') = 1,$$

be the factorization. Then

$$(V^r) = (\pi') = \tilde{\mathfrak{P}}^{a'}(\tilde{\mathfrak{P}}')^a\delta',$$

and

$$(p^r) = (F^rV^r) = (\pi\pi') = (\tilde{\mathfrak{P}}\tilde{\mathfrak{P}}')^{a+a'}\delta\delta'.$$

Then (6) yields $r = a+a'$. Suppose that $G_{\tilde{\mathfrak{P}}} = G_{1.0}$. Since V is an isomorphism on $\mathbb{M}(G_{\tilde{\mathfrak{P}}})$ so is π'. But by definition the action of $\mathcal{O}_K$ on $\mathbb{M}(G_{\tilde{\mathfrak{P}}})$ (resp. $\mathbb{M}(G_{\tilde{\mathfrak{P}}'})$) factors through an imbedding $\mathcal{O}_K\hookrightarrow\mathcal{O}_{K,\mathfrak{P}}\simeq\mathbb{Z}_p$ (resp. $\mathcal{O}_K\hookrightarrow\mathcal{O}_{K,\mathfrak{P}'}\simeq\mathbb{Z}_p$), π' should be a unit in $\mathcal{O}_{K,\mathfrak{P}}$ and $a' = 0$. This shows that $(V^r) = (\tilde{\mathfrak{P}}')^r\delta'$ and $(F^r) = \tilde{\mathfrak{P}}^r\delta$. Thus F^r is an isomorphism on $\mathbb{M}(G_{\tilde{\mathfrak{P}}'})$, which implies that $G_{\tilde{\mathfrak{P}}'}\simeq G_{0.1}$ and $G_{\mathfrak{P}}\simeq G_{1.0}\times G_{0.1} = \mathbb{L}$. If $G_{\tilde{\mathfrak{P}}} = G_{0.1}$ the similar argument will show the claim.

□

Lemma 3.2. *For $\mathfrak{P} \in \mathcal{P}_{inert}$, $G_{\mathfrak{P}}$ is isogeneous to $G_{1.1}^{f(\mathfrak{P}/p)}$.*

Proof. By **Fact 2.1** $G_{\mathfrak{P}}$ is isogeneous to $\prod G_{d.c}$. We have to show all simple factors are isomorphic to $G_{1.1}$. We will use the charcterization (4). Let us take a positive integer r as **Lemma 3.1** and

$$(F^r) = (\pi) = \mathfrak{P}^a \delta, \quad (\mathfrak{P}, \delta) = 1,$$

be the factorization. Since $\mathfrak{P}' = \mathfrak{P}$,

$$(V^r) = (\pi') = \mathfrak{P}^a \delta', \quad (\mathfrak{P}, \delta') = 1,$$

and we obtain

$$G_{\mathfrak{P}}[F^r] = G_{\mathfrak{P}}[V^r]. \tag{10}$$

Let G be a simple factor of $G_{\mathfrak{P}}$. Then (10) yields

$$G[F^r] = G[V^r],$$

and $G = G_{1.1}$ by (4).

□

Put $\mathcal{P} = \{\mathfrak{P}_1, \cdots, \mathfrak{P}_t\}$ and the following proposition yields **Theorem 1.1** and **Theorem 1.2**. For a finite set S we denote the cardinality by $|S|$.

Proposition 3.1. (1) *Suppose $\mathcal{P} = \mathcal{P}_{inert}$. Then A is supersingular.*

(2) *The a-number of A is greater than or equal to $|\mathcal{P}_{inert}^{f=1}|$.*

(3) *Suppose that*

$$\mathcal{P} = \mathcal{P}_{inert}^{f=1} \cup \mathcal{P}_{split}^{f=1}.$$

Then

$$a(A) = |\mathcal{P}_{inert}^{f=1}|, \quad f(A) = |\mathcal{P}_{split}^{f=1}|.$$

In particular if $\mathcal{P} = \mathcal{P}_{inert}^{f=1}$ (resp. $\mathcal{P} = \mathcal{P}_{split}^{f=1}$) A is superspecial (resp. ordinary).

Proof. (1) is an immediate consequence of **Lemma 3.2** and **Fact 2.3**. Let $\{\mathfrak{P}_1, \cdots, \mathfrak{P}_s\}$ be the elements of $\mathcal{P}_{inert}^{f=1}$. Then $A[p^\infty]$ contains $G_{\mathfrak{P}_1} \times \cdots \times G_{\mathfrak{P}_s}$. By (8) we see that $h(G_{\mathfrak{P}_i}) = 2$ and **Lemma 3.2** implies that $G_{\mathfrak{P}_i}$ is isomorphic to $G_{1.1}$ for all i. Therefore $A[p]$ contains $G_{1.1}[p]^s$ and by (5),

$$a(A) = \dim_{\overline{\mathbb{F}}_p} \operatorname{Hom}(\alpha_p, A[p]) \geq \dim_{\overline{\mathbb{F}}_p} \operatorname{Hom}(\alpha_p, G_{1.1}[p]^s) = s.$$

Finally we will prove (3). Let us set $s = |\mathcal{P}_{inert}^{f=1}|$ and $g - s = |\mathcal{P}_{split}^{f=1}|$. Then **Lemma 3.1** and **Lemma 3.2** show that the product (7) becomes

$$A[p^\infty] \simeq G_{1.1}^s \times \mathbb{L}^{g-s},$$

and the claim follows.

□

Corollary 3.1. *Suppose that K is a Galois extension of $\mathbb{Q}$.*

(1) *If $\mathcal{P}_{inert}$ is not empty. Then A is supersingular.*

(2) *Suppose that* p *completely splits in* K_0. *If it also completely splits in* K A *is ordinary. Otherwise* A *is superspecial.*

4. Rationality

Let $\mathcal{A}$ be an abelian scheme over $\mathbb{Z}_p$ of relative dimension g acted by $\mathcal{O}_K$ where K is a CM-field so that $[K:\mathbb{Q}]=2g$ (Here we do *not* assume that the action is defined over $\mathbb{Z}_p$). Then there is an isomorphism

$$H^1_{DR}(\mathcal{A}/\mathbb{Z}_p)\simeq \mathcal{O}_K\otimes_{\mathbb{Z}}\mathbb{Z}_p, \tag{11}$$

and we fix one of them. In fact here is an example of such an isomorphism. Take a finite field extension k of $\mathbb{F}_p$ so that $\mathcal{O}_K$ is defined over $W(k)$ and there is an isomorphism $H^1_{DR}(\mathcal{A}/W(k))\simeq \mathcal{O}_K\otimes_{\mathbb{Z}} W(k)$ which can be written,

$$H^1_{DR}(\mathcal{A}/\mathbb{Z}_p)\otimes_{\mathbb{Z}_p} W(k)\simeq (\mathcal{O}_K\otimes_{\mathbb{Z}}\mathbb{Z}_p)\otimes_{\mathbb{Z}_p} W(k),$$

compatible with action of $\sigma\in \mathrm{Aut}_{\mathbb{Z}_p}W(k)$. Taking the invariant part by σ and we obtain the desired isomorphism. We assume that p completely splits in the totally real subfield K_0:

$$p=\mathfrak{P}_1\cdots\mathfrak{P}_g,$$

and that every prime $\mathfrak{P}_i$ remains prime in K for all i. Then $\mathcal{O}_K\otimes_{\mathbb{Z}}\mathbb{Z}_p\simeq W(\mathbb{F}_{p^2})^g$ and let e_i be the $\mathbb{Z}_p$-endomorphism of $H^1_{DR}(\mathcal{A}/\mathbb{Z}_p)$ which corresponds to the projection to the i-th factor via (11). Let A be the special fiber which is an abelian variety of dimension g over $\mathbb{F}_p$. Identifying $H^1_{DR}(\mathcal{A}/\mathbb{Z}_p)$ with $T_p(A)$ by **Fact 2.6** we think e_i to be an endomorphism of $T_p(A)$.

Lemma 4.1. *For every* i, e_i *is contained in* $\mathrm{End}_{\mathcal{D}(\mathbb{F}_p)}(T_p(A))$.

Proof. Let k be a finite extension of $\mathbb{F}_p$ over which $\mathcal{O}_K$ is defined and consider a Cartesian diagram:

$$\begin{tikzcd}
\mathrm{End}_{\mathcal{D}(\mathbb{F}_p)}(T_p(A)) \arrow[r] \arrow[d] & \mathrm{End}_{\mathbb{Z}_p}(T_p(A)) \arrow[d] \\
\mathrm{End}_{\mathcal{D}(k)}(T_p(A\otimes_{\mathbb{F}_p} k)) \arrow[r] & \mathrm{End}_{W(k)}(T_p(A\otimes_{\mathbb{F}_p} k)),
\end{tikzcd}$$

where the horizontal (resp. the vertical) arrows are the natural inclusions (resp. extensions over $W(k)$). Regarding e_i as an element of $\mathrm{End}_{W(k)}(T_p(A\otimes_{\mathbb{F}_p} k))$ it is sufficient to show that it is contained in $\mathrm{End}_{\mathcal{D}(k)}(T_p(A\otimes_{\mathbb{F}_p} k))$. But this is clear since it is the image of $(0,\cdots,0,1,0,\cdots,0)\in W(\mathbb{F}_{p^2})^g$ (where 1 places in the i-th position from the left) by the composition of

$$W(\mathbb{F}_{p^2})^g\simeq \mathcal{O}_K\otimes_{\mathbb{Z}}\mathbb{Z}_p\subset \mathrm{End}_k(A\otimes_{\mathbb{F}_p} k)\otimes_{\mathbb{Z}}\mathbb{Z}_p\overset{\tau}{\simeq}\mathrm{End}_{\mathcal{D}(k)}(T_p(A\otimes_{\mathbb{F}_p} k)),$$

where τ is the isomorphism of **Fact 2.5**.

□

Via the isomorphism of **Fact 2.5**,

$$\mathrm{End}_{\mathbb{F}_p}(A) \otimes_{\mathbb{Z}} \mathbb{Z}_p \overset{\tau}{\simeq} \mathrm{End}_{\mathcal{D}(\mathbb{F}_p)}(T_p(A)),$$

let $\epsilon_i \in \mathrm{End}_{\mathbb{F}_p}(A) \otimes_{\mathbb{Z}} \mathbb{Z}_p$ correspond to $e_i \in \mathrm{End}_{\mathcal{D}(\mathbb{F}_p)}(T_p(A))$. Since $\mathrm{End}_{\mathbb{F}_p}(A) \otimes_{\mathbb{Z}} \mathbb{Z}_p$ acts on $A[p^\infty]$ we set

$$G_i := \epsilon_i A[p^\infty].$$

which is a p-divisible group over $\mathbb{F}_p$. Thus we have a decomposition over $\mathbb{F}_p$:

$$A[p^\infty] = G_1 \times \cdots \times G_g.$$

Lemma 4.2. *The base change of G_i over $\overline{\mathbb{F}}_p$ is isomorphic to $G_{1,1}$ for all i.*

Proof. The construction shows

$$\mathbb{M}(G_i) = e_i H^1_{DR}(\mathcal{A}/\mathbb{Z}_p) = W(\mathbb{F}_{p^2}).$$

Therefore the height of G_i is 2 and **Proposition 3.1** (in particular its proof) implies the claim.

□

Proposition 4.1. *Suppose that p is greater than 3. Let J be an abelian variety of dimension g over $\mathbb{F}_p$ whose p-divisible group $J[p^\infty]$ is a product*

$$J[p^\infty] = G_1 \times \cdots \times G_g,$$

over $\mathbb{F}_p$ so that every base change of G_i over $\overline{\mathbb{F}}_p$ is isomorphic to $G_{1,1}$. Then it is a product of supersingular elliptic curve $\{E_i\}_{1\le i\le g}$

$$J = E_1 \times \cdots \times E_g,$$

defined over $\mathbb{F}_p$. In particular $\Phi_{J,p}(t) = (t^2+p)^g$.

Proof. The assumption and (5) imply that the a-number of J is g. By **Fact 2.4** we see that $J \otimes_{\mathbb{F}_p} \overline{\mathbb{F}}_p$ is isomorphic to a product of supersingular elliptic curves, $E_1 \times \cdots \times E_g$. A simple consideration shows that, if necessary changing indices, we may assume that $G_i = E_i[p^\infty]$. Let ϕ be the p-power Frobenius. Since J is defined over $\mathbb{F}_p$ we have a diagram:

$$\begin{tikzcd} E_i \arrow[r, "\phi"] \arrow[d, "\nu_i"'] & E_i^{\phi} \arrow[d, "\nu_i^{\phi}"'] \\ J \arrow[r, "\phi"] & J, \end{tikzcd}$$

where ν_i is the inclusion and it induces

$$\begin{tikzcd} E_i[p^\infty] \arrow[r, "\phi"] \arrow[d, "{\nu_i[p^\infty]}"'] & E_i^{\phi}[p^\infty] \arrow[d, "{\nu_i^{\phi}[p^\infty]}"'] \\ J[p^\infty] \arrow[r, "\phi"] & J[p^\infty]. \end{tikzcd}$$

By the assumption all $\{E_i[p^\infty](=G_i)\}_i$ and $\{\nu_i[p^\infty]\}_i$ are defined over $\mathbb{F}_p$ and we find that $E_i[p^\infty]=E_i^\phi[p^\infty]$ and that $\nu_i[p^\infty]=\nu_i^\phi[p^\infty]$ $(\forall i)$. Therefore $E_i=E_i^\phi$ and $\nu_i=\nu_i^\phi$, which imply that $\{E_i\}_i$ are defined over $\mathbb{F}_p$ and so is the product. The last claim follows from the well-known fact that the characteristic polynomial of p-power Frobenius of a supersingular elliptic curve over $\mathbb{F}_p$ is t^2+p if $p\geq 5$.

□

Now **Theorem 1.3** follows from **Lemma 4.2** and **Proposition 4.1**.

Proof of Corollaries. We will first show **Corollary 1.1**. Notice that K_0 is $\mathbb{Q}(\zeta_N+\zeta_N^{-1})$ and consider a sequence

$$1\to \mathrm{Gal}(K/K_0)\simeq\{\pm1\}\to\mathrm{Gal}(K/\mathbb{Q})\simeq(\mathbb{Z}/(N))^\times\to\mathrm{Gal}(K_0/\mathbb{Q})\to 1.$$

Then we see that $p\equiv -1(N)$ is equivalent to the fact that p completely splits in K_0 and each prime factor of p in K_0 remains prime in K. On the other hand $p\equiv 1(N)$ if and only if p completely splits in K. More generally an existence of a positive integer h so that $p^h\equiv -1(N)$ implies that an every prime factor of p in K_0 remains prime in K. Now the desired claims follow from **Theorem 1.4**. In the case of **Corollary 1.2** observe $K_0=\mathbb{Q}(\zeta_M+\zeta_M^{-1})$ and we find

$$\mathrm{Gal}(K/\mathbb{Q})\simeq\mathrm{Gal}(\mathbb{Q}(\zeta_M+\zeta_M^{-1})/\mathbb{Q})\times\mathrm{Gal}(\mathbb{Q}(\zeta_d)/\mathbb{Q}),$$

which is isomorphic to $((\mathbb{Z}/(M))^\times/\pm1)\times\{\pm1\}$. Now the assertion is clear.

□

5. EXAMPLES

Example 4.1. Let l be an odd prime and we define a curve $C(l)$ to be the smooth projective model of

$$y^l=x(1-x),$$

over $\mathbb{Q}$, which has a genus $(l-1)/2$. An l-th primitive root of unity ζ_l acts by

$$\zeta_l(x)=x,\quad \zeta_l(y)=\zeta_l y,$$

and $\mathbb{Z}[\zeta_l]$ is the endomorphism ring of $\mathrm{Jac}(C(l))$. Thus $C(l)$ is a CM-curve. Let p be a good prime which l divides p^h+1 for a certain positive integer h. Set $q=p^h$ and let F_{q+1} be a smooth projective model of a Fermat curve

$$X^{q+1}+Y^{q+1}=1,$$

over $\mathbb{Q}$. Then by

$$x=X^{q+1},\quad y=(XY)^{\frac{q+1}{l}},$$

we have a $\mathbb{Q}$-rational proper surjective morphism,

$$\pi: F_{q+1}\to C(l),$$

and $H^1_{et}(C(l)_p \otimes_{\mathbb{F}_p} \overline{\mathbb{F}}_p, \mathbb{Q}_l)$ is a Galois submodule of $H^1_{et}((F_{q+1})_p \otimes_{\mathbb{F}_p} \overline{\mathbb{F}}_p, \mathbb{Q}_l)$. It is known that $|F_{q+1}(\mathbb{F}_{p^2})|$ attains the Hasse-Weil upper bound $1 + p^2 + 2gp$ and

$$\Phi_{F_{q+1},p}(t) = (t^2 + p)^g,$$

where $g = q(q-1)/2$ is the genus of F_{q+1} ([10]**Example 6.3.6**). Thus we see that $\Phi_{C(l),p}(t) = (t^2 + p)^{(l-1)/2}$ and that

$$|C(l)_p(\mathbb{F}_{p^2})| = 1 + p^2 + (l-1)p,$$

if there is a positive integer h such that $p^h \equiv -1(l)$. In fact **Corollary 1.1** is weaker than this. Namely if $h = 1$ it recovers these results but if otherwise it only says that $C(l)$ has supersingular reduction. Therefore we find that **Corollary 1.1**(3) is merely a sufficient condition for a prime p at which the reduction of a CM-curve to be $\mathbb{F}_{p^2}$-maximal, i.e. the number of $\mathbb{F}_{p^2}$-rational points attains the Hasse-Weil upper bound.

Example 4.2.([3]**Example 2.3**) Let us consider a curve,

$$Y : y^3 = x(x^3 - 14x^2 - 49x - 49),$$

which is a smooth model of a Picard curve of CM-type. Its genus is 3 and the endomorphism ring of its Jacobian is $\mathbb{Z}[\zeta_7 + \zeta_7^{-1}, \zeta_3]$. Thus by **Corollary 1.2** we see that a good prime p is supersingular if $p \equiv -1(3)$. Moreover it is superspecial if $p \equiv \pm 1(7)$ and $p \equiv -1(3)$. Using the computation of [3] one can determine $\Phi_{Y,p}(t)$. In fact let C be a curve over $\mathbb{Q}$ defined by

$$C : y^3 = x(x^7 + 1).$$

Then Geemen, Koike and Weng have shown that $\mathrm{Jac}(C)$ is isogeneous to the product $\mathrm{Jac}(Y)^2 \times E$ with a CM-elliptic curve E. Moreover

Fact 5.1. *(*[3]**Corollary 3.1** *and* **Corollary 3.2***)*

1. *For* $p \equiv 2, 5, 11, 17(21)$,
$$\Phi_{C,p}(t) = (t^4 - pt^2 + p^2)^2(t^2 + p)^3.$$
2. *For* $p \equiv 8, 20(21)$,
$$\Phi_{C,p}(t) = (t^2 + p)^7.$$

Since Y has genus 3 the degree of $\Phi_{Y,p}(t)$ should be six and we find

- For $p \equiv 2, 5, 11, 17(21)$,
$$\Phi_{Y,p}(t) = (t^4 - pt^2 + p^2)(t^2 + p),$$
- For $p \equiv 8, 20(21)$,
$$\Phi_{Y,p}(t) = (t^2 + p)^3.$$

These coincide with the results of **Corollary 1.2** (see also comments after **Theorem 1.1**). Moreover in the second case we know that $\mathrm{Jac}(Y_p)$ is a product of supersingular elliptic curves over $\mathbb{F}_p$ and that $|Y_p(\mathbb{F}_{p^2})| = 1 + p^2 + 6p$.

Example 4.3.([3]**Example 2.1**) Remember that the *n-th Chebyshev polynomial* U_n is defined by a recursive relation:

$$U_{n+1}(x) = xU_n(x) - U_{n-1}(x), \quad U_0(x) = 2, \quad U_1(x) = x.$$

Using this for a prime $l \geq 5$ we define a curve X_l defined over $\mathbb{Q}$ by

$$X(l) : y^2 = U_l(x),$$

which is a quotient of a hyperelliptic curve

$$y^2 = x(x^{2l} + 1),$$

by an involution

$$\tau : (x, y) \mapsto (\frac{1}{x}, \frac{y}{x^{l+1}}).$$

It has a genus $(l-1)/2$ and the endomorphism ring of $\mathrm{Jac}(X(l))$ is $\mathbb{Z}[\zeta_l + \zeta_l^{-1}, \zeta_4]$. Thus $X(l)$ is a CM-curve. **Corollary 1.2** shows that a good prime p satisfying $p \equiv -1(4)$ is supersingular. Moreover if $p \equiv -1(4)$ and $p \equiv \pm 1(l)$, $\mathrm{Jac}(X(l)_p)$ is a product of supersingular elliptic curves over $\mathbb{F}_p$ and $|X(l)_p(\mathbb{F}_{p^2})|$ attains the Hasse-Weil upper bound $1 + p^2 + (l-1)p$.

REFERENCES

[1] P. Deligne, Variétés abéliennes ordineire sur un corps fini, . *Inventiones Math*, 2:238–243, 1967.
[2] M. Demazure, Lectures on p-divisible groups. *Lecture Notes in Mathematics*, Springer-Verlag, 1972.
[3] B. van Geemen, K. Koike and A. Weng, Quotient of Fermat curves and a Hecke character. *Finite Fields and Their Appl.*, 11:6–29, 2005.
[4] A. Grothendieck, Groupes de Barsotti-Tate et cristaux de Diuedonné. *Sém. Math. Sup. 45*, Presses de l'Univ. de Montreal, 1970.
[5] H.W. Lenstra and F. Oort, Simple abelian varieties having a prescribed formal isogeny type. *J. Pure Appl. Algebra.*, 4:47–53, 1974.
[6] Yu.I. Manin, The theory of commutative formal groups over fields of finite characteristic. *Russ. Math. Surv.*, 18:1-80, 1963.
[7] J.S. Milne and W.C. Waterhaus, Abelian varieties over finite fields. *Proc. Symp. Pure Math.*, 20:53–64, 1971.
[8] F. Oort, Abelian varieties over finite fields. *Lecture note of Summer school in Göttingen*, June, 2007.
[9] F. Oort, Which abelian surfaces are products of elliptic curves. *Math. Ann.*, 214:35–47, 1975.
[10] H. Stichtenoth, Algebraic function fields and codes. *Springer-Verlag, Berlin*, 1993.
[11] J. Tate, Endomorphism of abelian varieties over finite fields. *Inventiones Math*, 2:134–144, 1966.
[12] W.C. Waterhouse, Abelian varieties over finite fields. *Ann. Sc. Ec. Norm. Sup.*, 4-2:521–560, 1969.
[13] A. Zaytsev, Generalization of Deuring reduction theorem. *arXiv [math.AG]*, 1209.5207v1.